University Health™	Policy #: Rad Proc 13.14.10
SUBJECT: CT Pancreas 6.10	Effective: 10/2013 Reviewed: 03/07/2017
APPROVED BY: Body Imaging Medical Director	Page 1 of 3

Purpose: To provide computed tomography staff with the required protocol for performing CT

**Pancreas** 

Scope: All adult patients 18 years and older.

**Clinical Indication: Acute or chronic pancreatitis** 

Patient Preparation: Clear liquid diet (6 hours prior to exam)

Orientation: Feet first Breathing: Inspiration

Oral Contrast: VoLumen Protocol (see proc. 6.12.1)

IV Contrast per Weight: 1ml/lb or 2ml/kg not to exceed 150ml injected @ 4ml/sec

Coverage: Dome of diaphragm through iliac crest; for pelvis continue through ischial tuberosities

Anatomic Reference: Xiphoid process; Scan Delay: Smart prep or care bolus

**Group 1: Non-contrast – Dome of diaphragm to iliac crest** 

Scan Mode	Thickness Speed Pitch	Table Interval	SFOV	kVp	Auto mA/ Noise Index	Prep Time (sec)	Recon Type
LS 16 0.8 sec Helical Full	2.5 27.50 1.375:1	2.5	Large	120	80-440 11.5	N	Standard
VCT 0.8 sec Helical Full	2.5 55.00 1.375:1	2.5	Large	120	80-600 11.5	N	Standard
AS 64 0.5 sec Helical	3 1	3	360	120	Ref MAS 250	N	Standard

Group 2: Arterial Phase – Dome of diaphragm to iliac crest

Scan Mode	Thickness Speed Pitch	Table Interval	SFOV	kVp	mA Auto mA/ Noise Index	Prep Time (sec)	Recon Type
LS 16 0.8 sec Helical	2.5 27.50 1.375:1	2.5	Large	120	80-440 11.5	30 sec.	Standard
VCT 64 0.8 sec Helical Full	2.5 55.00 1.375:1	2.5	Large	120	80-600 11.5	30 sec	Standard
AS 64 0.5 sec	3 1	3	380	120	Ref MAS 250	30 sec	Standard

**Group 3:** Venous Phase- Dome of diaphragm to iliac crest

Scan Mode	Thickness Speed Pitch	Table Interval	SFOV	kVp	mA Auto mA/ Noise Index	Prep Time (sec)	Recon Type
LS 16 0.8 sec Helical	2.5 27.50 1.375:1	2.5	Large	120	80-440 11.5	40 sec	Standard
VCT 64 0.8 sec Helical Full	2.5 55.00 1.375:1	2.5	Large	120	80-600 11.5	40 sec	Standard
AS 64 0.5 sec	3 1	3	380	120	Ref MAS 250	40 sec	Standard

University Health™	Policy #: Rad Proc 13.14.10
SUBJECT: CT Pancreas 6.10	Effective: 10/2013 Reviewed: 03/07/2017
APPROVED BY: Body Imaging Medical Director	Page 3 of 3

**Algorithm: Recon 1 & 2 Standard** 

**Reformation: 2.0 Sagittal and Coronal reformations of the venous phase** 

Network: Recon 1 (5mm) to PACS. Recon 2 (1.25) auto transmit to AWSERV

Notes: Auto mA must be on and mA table checked prior to scanning.