

 University Health™	Policy #: Rad Proc 13.14.18
SUBJECT: CT Abdomen/Pelvis Endostent Graft Follow-up	Effective: 10/2013 Reviewed: 03/07/2017
APPROVED BY: Body Imaging Medical Director	Page 1 of 3

Purpose: To provide computed tomography staff with the required protocol for performing CT Endostent Graft Follow-up

Scope: All adult patients 18 years and older.

Clinical Indication: Follow up to Endostent Graft

Patient Preparation: Clear liquid diet (6 hours prior to exam)

Orientation: Feet first

Breathing: Inspiration

Oral Contrast: VoLumen Protocol (see proc. 6.12.1)

IV Contrast per Weight: 1ml/lb or 2ml/kg not to exceed 150ml injected @ 4ml/sec

Coverage: Dome of diaphragm through ischial tuberosities

Anatomic Reference: Xiphoid process;

Scan Delay: Smart prep or care bolus

Group 1: Non-contrast – Dome of diaphragm to ischial tuberosities

Scan Mode	Thickness Speed Pitch	Table Interval	SFOV	kVp	Auto mA/ Noise Index	Prep Time (sec)	Recon Type
LS 16 0.8 sec Helical Full	5 27.50 1.375:1	5	Large	120	80-440 11.5	N	Standard
VCT 0.8 sec Helical Full	5 55.00 1.375:1	5	Large	120	80-600 11.5	N	Standard
AS 64 0.5 sec Helical	5 1	5	360	120	Ref MAS 250	N	Standard

 University Health™	Policy #: Rad Proc 13.14.18
SUBJECT: CT Abdomen/Pelvis Endostent Graft Follow-up	Effective: 10/2013 Reviewed: 03/07/2017
APPROVED BY: Body Imaging Medical Director	Page 2 of 3

Group 2: Arterial Phase – Dome of diaphragm to Ischial tuberosities

Scan Mode	Thickness Speed Pitch	Table Interval	SFOV	kVp	mA Auto mA/ Noise Index	Prep Time (sec)	Recon Type
LS 16 0.8 sec Helical	2.5 27.50 1.375:1	2.5	Large	120	80-440 11.5	Smart Prep Aorta Near Stent	Standard
VCT 64 0.8 sec Helical Full	2.5 55.00 1.375:1	2.5	Large	120	80-600 11.5	Smart Prep Aorta Near Stent	Standard
AS 64 0.5 sec	3 1	3	380	120	Ref MAS 250	Care Bolus Aorta Near Stent	Standard

Group 3: Delayed Phase – Dome of diaphragm to Ischial tuberosities

Scan Mode	Thickness Speed Pitch	Table Interval	SFOV	kVp	mA Auto mA/ Noise Index	Prep Time (sec)	Recon Type
LS 16 0.8 sec Helical	5 27.50 1.375:1	5	Large	120	80-440 11.5	120 sec	Standard
VCT 64 0.8 sec Helical Full	5 55.00 1.375:1	5	Large	120	80-600 11.5	120 sec	Standard
AS 64 0.5 sec	5 1	5	380	120	Ref MAS 250	120 sec	Standard

 University Health™	Policy #: Rad Proc 13.14.18
SUBJECT: CT Abdomen/Pelvis Endostent Graft Follow-up	Effective: 10/2013 Reviewed: 03/07/2017
APPROVED BY: Body Imaging Medical Director	Page 3 of 3

Algorithm: Recon 1 & 2 Standard

Reformation: Sagittal and Coronal reformations of the delayed phase.

Network: Recon 1 (5mm) to PACS. Recon 2 (1.25) auto transmit to AWSERV

Notes: Auto mA must be on and mA table checked prior to scanning.